

Statistical Predictions From Anarchic Field Theory Landscapes

Vijay Balasubramanian^{1,ab}, Jan de Boer^{2,c}, and Asad Naqvi^{3,de},

^a *David Rittenhouse Laboratories, University of Pennsylvania, Philadelphia, PA 19104, USA*

^b *School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA*

^c *Instituut voor Theoretische Fysica, Valckenierstraat 65, 1018XE Amsterdam, The Netherlands*

^d *Department of Physics, Swansea University, Singleton Park, Swansea, SA2 8PP, UK*

^e *Department of Physics, Lahore University of Management and Sciences, Lahore, Pakistan*

Abstract

Consistent coupling of effective field theories with a quantum theory of gravity appears to require bounds on the rank of the gauge group and the amount of matter. We consider landscapes of field theories subject to such boundedness constraints. We argue that appropriately “coarse-grained” aspects of the randomly chosen field theory in such landscapes, such as the fraction of gauge groups with ranks in a given range, can be statistically predictable. To illustrate our point we show how the uniform measures on simple classes of $\mathcal{N} = 1$ quiver gauge theories localize in the vicinity of theories with certain typical structures. Generically, this approach would predict a high energy theory with very many gauge factors, with the high rank factors largely decoupled from the low rank factors if we require asymptotic freedom for the latter.

¹e-mail: vijay@physics.upenn.edu

²e-mail: J.deBoer@uva.nl

³e-mail: a.naqvi@swansea.ac.uk

Contents

1	Introduction	1
2	The set of field theories	2
2.1	Interesting classes of quiver gauge theories	3
2.2	Averages and typicality	4
2.3	Choice of measure	4
3	Typicality in Toy Landscapes	6
3.1	Theories without matter: coarse graining and typicality	6
3.2	Cyclic, chiral quivers	7
3.2.1	The canonical ensemble breaks down	8
3.2.2	Microcanonical analysis	11
4	Thinking about the general quiver	13
4.1	Implementing anomaly cancellation	14
4.2	Dealing with discrete quiver symmetries: an example	16
4.3	Towards dynamics	18
4.3.1	Four node, asymptotically free quivers	18
4.3.2	General quiver with unequal gauge groups	18
5	Conclusion	19

1 Introduction

It is commonly supposed that the huge numbers of vacua that can arise from different compactifications of string theory [1, 2] imply a complete loss of predictability of low energy physics. If this is the case, the stringiness simply constrains the possible dynamics rather than the precise complement of forces and matter. Every string theory leads to some effective field theory at a high scale Λ , taken to be, say, an order of magnitude below the string scale. Predictions for low energy physics have to be made in terms of this effective field theory. Thus, the landscape of string theory vacua leads to a landscape of effective field theories at the scale Λ . Here we ask if constraints of finiteness imposed on this landscape via its origin in string theory might be sufficient to lead to a degree of predictability, at least in some statistical sense. Previous authors have discussed how continuous parameters can scan in a random landscape of effective field theories [3, 4, 5, 6, 7, 8, 9], and there has been some study of the gauge groups and matter content attainable from specific string theoretic scenarios [10, 11, 12, 13, 14, 15]. For example, [14] and [15] discuss the distribution of gauge groups arising in intersecting brane models on torus orientifolds.

We will impose the weakest of the constraints arising from string theory – namely that it should be possible to couple the effective field theory consistently to a quantum theory of gravity. It has been argued [16, 17, 18] that such consistency with string theory requires that the rank of the gauge group and the number of matter fields

be bounded from above.¹ Since we will not impose any constraints based on rules arising from symmetry or dynamics on the measure, we will call this an “anarchic” landscape, in recollection of the terminology in [6]. Thus we will study simple anarchic landscapes of field theories bounded in this way, and illustrate how statistics can lead to characteristic predictions for the low energy physics. These predictions are strongest for appropriately coarse-grained attributes of a theory that possess the property of *typicality* in such landscapes – i.e. they are overwhelmingly likely to lie close to certain typical values. An example of such a typical property will be the fraction of gauge groups with ranks lying within some range. We will illustrate and develop our thinking using some simple examples.

2 The set of field theories

A natural, large class of field theories to consider is the set of quiver gauge theories. For simplicity, we will restrict attention to $\mathcal{N} = 1$ supersymmetric quiver gauge theories where the gauge group is a product of unitary groups,

$$G = \prod_{i=1}^L U(N_i). \quad (1)$$

In addition, there will be A_{ii} hypermultiplets transforming in the adjoint of $U(N_i)$, and A_{ij} hypermultiplets transforming in the $(\mathbf{N}_i, \bar{\mathbf{N}}_j)$ of $U(N_i) \times U(N_j)$. The nonnegative integer matrix

$$A_{ij} \geq 0, \quad i, j = 1 \dots L \quad (2)$$

describes the number of arrows from site i to site j in the quiver.

Of course, to specify the full gauge theory we also need to describe the Kähler potential for the hypermultiplets, the gauge kinetic terms, the superpotential and possibly Fayet-Iliopoulos terms. We will postpone a discussion of these quantities for now and will in this paper only discuss the matter and gauge group content of the $\mathcal{N} = 1$ theory.

Gauge theories of quiver type are ubiquitous in string theory, and this is the main motivation to restrict attention to this class. Bifundamentals tend to appear in string theory because strings have two endpoints only. A typical setup to engineer quiver $\mathcal{N} = 1$ theories is to consider D6-branes wrapping 3-cycles inside a Calabi-Yau manifold in type IIA string theory, in which case the number of bifundamentals is related to the intersection number of the 3-cycles. By including orientifolds, we can also easily engineer quiver gauge theories with SO and Sp gauge factors, but we will postpone a study of these theories to another occasion.

Our goal will thus be to study random $U(N)$ quiver gauge theories. Before looking at some concrete examples, we are first going to make some general remarks on possible further restrictions on the set of gauge theories, on the choice of measure on the space of theories, and the kinds of properties we might predict.

¹A possible bound on the number of matter species in theories containing gravity was originally discussed by Bekenstein [19].

2.1 Interesting classes of quiver gauge theories

Three possible restricted sets of gauge theories are:

1. Anomaly free theories. A simple physical requirement that we should impose is the condition that there are no anomalies. This translates into the statement that for all i

$$\sum_{j \neq i} (A_{ij} - A_{ji}) N_j = 0. \quad (3)$$

The expectation value of the left hand side of this equation in the unconstrained set of quiver gauge theories with the uniform measure is clearly zero, as the measure is invariant under $A_{ij} \leftrightarrow A_{ji}$. Therefore, “on average,” random quiver gauge theories are anomaly free, and one might be inclined to not worry about anomalies anymore. However, from a physical point of view it seems rather peculiar to allow forbidden theories in an ensemble, as it is not at all clear that properties of the set of anomaly free theories are properly reproduced by the full set of random quiver gauge theories. Hence we will for the most part restrict to field theories which obey (3).

2. Asymptotically free theories. Another natural constraint we can impose is that the theories we consider are asymptotically free, which makes them well-defined in the UV. Asymptotic freedom is less compelling than anomaly cancellation, as the set of random quiver theories may well represent a set of low-energy effective field theories obtained e.g. in string theory. Gauge group factors that are IR free and strongly coupled in the UV will typically start to act as global symmetries at sufficiently low energies and will not directly lead to contradictions. The condition for asymptotic freedom is that for all i ,

$$A_{ii} N_i + \sum_{j \neq i} (A_{ij} + A_{ji}) N_j < 3 N_i. \quad (4)$$

This tends to constrain the A_{ij} to not be very large but to be of order unity instead.

3. Purely chiral theories. If we imagine our field theory to be some effective field theory at a high scale M , assume there are no other dimensionful parameters around, and write down the most general superpotential, it will contain many mass terms with masses of order M . At energies below M , it makes sense to integrate out all massive fields with masses of order M . The remaining gauge theory will have fewer fields and will no longer allow for mass terms: all fields that can be integrated out have been removed. The remaining set of purely chiral theories with

$$A_{ii} = 0, \quad A_{ij} = 0 \text{ or } A_{ji} = 0 \text{ for } i \neq j \quad (5)$$

are therefore a natural starting point for viewing random quivers as low-energy effective field theories. Such chiral theories allow for general cubic superpotentials at the marginal level. Higher order terms are suppressed by a mass scale

in the Lagrangian, although some quartic superpotentials can become marginal in the infrared.

4. Equal rank theories. In order to simplify the analysis, we could take the ranks of all the gauge groups to be fixed and equal. For such theories both the anomaly cancellation constraint as well as the asymptotic freedom constraint are much easier to implement. However, we do not have an obvious physical motivation that would prefer these theories, so they are mainly helpful to develop intuition for the more general case.

2.2 Averages and typicality

Given a set of gauge theories with a suitable measure on them, we can compute expectation values of various quantities, such as the average rank of a gauge group, the average number of matter fields, etc. Though averages are useful to compute, they are especially interesting when they also represent the *typical* value of a quantity. Typicality is a notion that exists in situations when a thermodynamic limit can be taken wherein some parameter N , controlling the size of the ensemble, can be taken to infinity. Then, a quantity enjoys the property of typicality if its probability distribution becomes more and more narrowly peaked around its expectation value as $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} \frac{\langle \mathcal{O}^2 \rangle - \langle \mathcal{O} \rangle^2}{\langle \mathcal{O} \rangle^2} = 0. \quad (6)$$

In other words, quantities that are typical are equal to their ensemble averages with probability one in the limit $N \rightarrow \infty$ ².

Familiar examples of typical operators are statistical mechanical quantities such as pressure and free energy. Also, we note that for a standard Boltzmann distribution, for one particular occupation number with

$$\langle N \rangle = \frac{\sum_{k \geq 0} k e^{-\beta k}}{\sum_{k \geq 0} e^{-\beta k}} = \frac{e^{-\beta}}{1 - e^{-\beta}}, \quad \langle N^2 \rangle = \frac{\sum_{k \geq 0} k^2 e^{-\beta k}}{\sum_{k \geq 0} e^{-\beta k}} = \frac{e^{-\beta}(1 + e^{-\beta})}{(1 - e^{-\beta})^2} \quad (7)$$

the variance to mean squared ratio appearing in (6) equals e^β . In other words, a microscopic quantity like a single occupation number will not be typical. Observables that achieve typicality are inevitably coarse-grained – e.g. the number of Boltzmann particles with energies between c/β and $(c + \epsilon)/\beta$ for constants c and ϵ will be typical. In studying the statistics of effective field theories we should be interested in finding appropriately “coarse-grained” structures that are typical.

2.3 Choice of measure

In order to define and discuss averages and typicality for random quiver gauge theories, we need to define a suitable measure on this space. One could imagine that

²This criterion is not very useful when $\langle \mathcal{O} \rangle = 0$. More generally, we should normalize the operator $\mathcal{O}$ in such a way that the range of values it can take is independent of N and then require that the variance vanishes in the large N limit.

dynamics gives rise to an interesting and complicated measure. For example, one could imagine weighing field theories by the dimension or even size of the cohomology of their respective moduli spaces, having the close connection between quiver gauge theories and D-brane moduli spaces in mind. As another simple example of how dynamics can affect the measure, if we suppose that dynamical effects can give the matter fields any expectation value, then generically all the gauge groups will be broken to $U(1)$ and analysis of the distribution of gauge factors is moot. However, in $N = 1$ theories of the kind we study, the potential for the matter fields typically develops isolated minima and the gauge group is broken to a product of Abelian and non-Abelian factors (for instance, a cubic superpotential for an adjoint superfield classically breaks $U(N) \rightarrow U(p) \times U(N-p)$ for some p). Classically, in the context of Calabi-Yau compactification, one imagines that the manifold has some set of distinct but intersecting cycles and the non-abelian factors in the gauge theory are related to the number of branes wrapped on each cycle. Then strong gauge dynamics might break these gauge factors further. For the present we will ignore such dynamical issues and use a uniform measure subject to various constraints of boundedness. Since we are ignoring rules arising from the underlying dynamics, we will call our measures “anarchic”.

Finally, in the context of e.g. string landscape discussions, one might want to associate various kinds of Bayesian measures to different types of field theories. For example, to correctly make statistical predictions for the UV field theory, given our hypothetical bound on the the matter and gauge groups, we strictly speaking condition our probability distribution on the known facts about infrared physics. From this perspective, we actually want the uniform measure on a bounded space of gauge theories that, when run to the infrared, contain the standard model as a sector. Conditioning in this way, is well beyond our ability at present, and so we will simply investigate the uniform measure on bounded spaces of quiver gauge theories, to study whether and how typicality occurs.

Experience in statistical physics has shown that directly computing averages and variances over bounded configuration spaces can be difficult. Thus, to simplify analysis we can try to use a grand canonical ensemble to constrain the total rank and the total number of matter fields. This involves summing over theories with arbitrary ranks and amounts of matter while including in the measure a Boltzmann factor for the rank of the gauge group, and a separate Boltzmann factor for the total number of matter fields

$$\rho \sim \exp(-\beta \sum_i N_i - \lambda \sum_{ij} A_{ij} N_i N_j). \quad (8)$$

One could also include Boltzmann factors for, e.g., the total number of nodes, the total number of gauge bosons, etc., but for our purposes (8) will be sufficient to illustrate the main ideas. Such an approach only works if the ensemble of theories does not grow exponentially fast in the total rank and number of matter fields. If such exponential growth occurs, the Boltzmann weight does not fall quickly enough for the microcanonical ensemble to be well approximated by the canonical ensemble. We will see that the space of theories typically grows too fast with the number of

fields to permit use of the canonical approach to make statistical predictions from a bounded landscape of effective field theories.

3 Typicality in Toy Landscapes

3.1 Theories without matter: coarse graining and typicality

As an example of our approach, consider a landscape of field theories with no matter, where the rank of the gauge group is equal to a large number N . For simplicity, let the gauge group be a product of unitary factors

$$G = \prod_{i=1}^L U(N_i). \quad (9)$$

Then the rank of G is $\sum_i N_i = N$; thus the N_i form an integer partition of N . To study the distribution of gauge factors in an anarchic landscape of such field theories, we can construct the canonical partition function

$$Z = \sum_{\{r_k\}} e^{-\beta \sum_k k r_k - \alpha \sum_k r_k} = \prod_k \frac{1}{1 - e^{-\beta k - \alpha}} \equiv \prod_k \frac{1}{1 - u q^k} \quad (10)$$

Here r_k is the number of gauge factors of rank k , β is a Lagrange multiplier constraining the total rank to be N and α is a Lagrange multiplier that can be used to constrain the number of gauge factors; sometimes it is more convenient to work with $q = e^{-\beta}$ and $u = e^{-\alpha}$ instead. In writing this we have used measure that treats the ordering of gauge factors as irrelevant. So, for example, $U(2) \times U(3) \times U(2)$ is the same as $U(3) \times U(2) \times U(2)$ and so on. In such a measure, all $U(N_i)$ factors are treated as identical, and not distinguished from each other by parameters like their gauge couplings. This measure will be modified if the gauge theory is realized by wrapping D-branes on cycles of Calabi-Yau because in that case the locations of branes and the sizes of the cycles will allow us to distinguish between many different configurations that lead to the same gauge group. Nevertheless, the present measure is interesting from a purely field theoretic point of view, i.e. if one is simply counting field theories, and is illustrative.

To fix β and α we require that

$$N = \sum_{j=1}^{\infty} \frac{j u q^j}{1 - u q^j} \quad ; \quad L = \sum_j \frac{u q^j}{1 - u q^j}, \quad (11)$$

where N is the total rank and L is the total number of gauge factors. We will take

$$u \sim O(1) \quad ; \quad \beta \sim \frac{1}{\sqrt{N}} \quad (12)$$

which, we will see later, implies $L \sim \sqrt{N}$. Then from (10) it is easy to show that:

$$\langle r_j \rangle = \frac{u q^j}{1 - u q^j} \quad ; \quad \text{Var}(r_j) = \frac{u q^j}{(1 - u q^j)^2} = \frac{\langle r_j \rangle}{1 - u q^j}. \quad (13)$$

The variance to mean squared ratio is

$$\frac{\text{Var}(r_j)}{\langle r_j \rangle^2} = \frac{1}{u q^j} = e^{\beta j + \alpha} \geq e^\alpha \geq O(1). \quad (14)$$

To get the last inequality we simply used $\alpha, \beta > 0$. Thus we see that in a universe with such anarchic landscapes, the number of gauge factors r_j with rank j is not typical in the sense defined in (6) and thus cannot be predicted with confidence.

However, we could ask whether there are any more coarse grained structures in such landscapes which are more predictable. For example, consider the number of gauge factors whose ranks lie between $c\sqrt{N}$ and $(c + \epsilon)\sqrt{N}$ where c and ϵ are both $O(1)$:

$$\langle R(c, \epsilon) \rangle \approx \int_{c\sqrt{N}}^{(c+\epsilon)\sqrt{N}} dj \langle r_j \rangle = \frac{1}{\beta} \ln \left[\frac{1 - u e^{-(c+\epsilon)\sqrt{N}\beta}}{1 - u e^{-c\sqrt{N}\beta}} \right], \quad (15)$$

where we approximated the sum as an integral. The variance of this coarse-grained variable is

$$\text{Var}(R(c, \epsilon)) = \int_{c\sqrt{N}}^{(c+\epsilon)\sqrt{N}} dj \text{Var}(r_j) = \frac{u}{\beta} \left[\frac{e^{-c\sqrt{N}\beta} - e^{-(c+\epsilon)\sqrt{N}\beta}}{(1 - u e^{-c\sqrt{N}\beta})(1 - u e^{-(c+\epsilon)\sqrt{N}\beta})} \right], \quad (16)$$

where we used the fact that in this canonical ensemble the r_j are statistically independent variables. Thus, for $\beta \sim 1/\sqrt{N}$ (12),

$$\langle R(c, \epsilon) \rangle \sim O(\sqrt{N}) \quad ; \quad \text{Var}(R(c, \epsilon)) \sim O(\sqrt{N}) \quad \implies \quad \frac{\text{Var}(R(c, \epsilon))}{\langle R(c, \epsilon) \rangle^2} \sim O(1/\sqrt{N}). \quad (17)$$

This means that the variance to mean squared ratio vanishes in the large N limit indicating that $R(c, \epsilon)$ is a typical variable. Thus, in such anarchic landscapes, the number of gauge factors with ranks between $c\sqrt{N}$ and $(c + \epsilon)\sqrt{N}$ can be predicted with high confidence. Also, approximating the second equation in (11) as an integral, the total number of gauge factors turns out to be

$$L = -\frac{\ln(1-u)}{u\beta} \sim O(\sqrt{N}). \quad (18)$$

By the above variance analysis this number will also be typical. Thus, in such anarchic landscapes, the total number of gauge factors is highly predictable. These results follow essentially because the unordered partitions of a large integer enjoy a sort of central limit theorem – representing such partitions by a Young diagram, one can show that in the large N limit, the boundary of an appropriately rescaled diagram approaches a limit shape encoded by the $\langle r_j \rangle$ computed above [20].

3.2 Cyclic, chiral quivers

Above we saw how suitably coarse-grained aspects of the structure of a randomly chosen field theory in a bounded landscape might be statistically predictable. The

next step is to add matter to the theory to see how this changes the analysis. As we discussed, we must insist that matter is added in an anomaly-free way and implementing this constraint is one of the main difficulties in studying statistical ensembles of quiver gauge theories. Thus, to make a beginning, we will study cyclic, chiral quiver gauge theories for which anomaly cancellation is very easy to implement.

In cyclic quivers, each gauge group is connected to the next one by bifundamentals, with the circle being completed when the last group connects to the first one. Taking the i th group around the circle to be $U(N_i)$, the constraint on the total rank will be $\sum_i N_i = N$. So, as in the example without matter, the number N_i form a partition of N . Anomaly cancellation requires that each gauge group have as many fundamentals as antifundamentals. It is easy to show that, the minimal solution to anomaly cancellation constraints is that the number of bifundamentals between $U(N_i)$ and $U(N_{i+1})$ is

$$A_{i(i+1)} = C^{-1} \cdot \prod_{l \neq i, (i+1)} N_l \quad ; \quad C = \text{GCD}(\{ \prod_{l \neq i, (i+1)} N_l \}) \quad (19)$$

All other solutions to the anomaly cancellation equations are integer multiples of (19). We will examine an ensemble in which the matter fields in the gauge theory are presumed to satisfy (19) in such a way that the total number of fields comes as close as possible to some bound K . Thus for this setup the matter fields are uniquely chosen once the gauge groups are selected. (More generally, we could have imagined an ensemble where the number of matter fields was allowed to vary, in which one would need to sum over multiples of $A_{i(i+1)}$ subject to a bound. This is difficult to do here since the GCD of the products of integer subsets appearing in the denominator of (19) is presumably sporadically behaved.)

One key difference from the matter-free case, is that the *order* in which the gauge groups appear on the ring of the quiver is important. In general, different orderings will lead to different quiver gauge theories, except when the permutations correspond to symmetries of the quiver, such as the cyclic permutations of the nodes, or cyclic permutations combined with one reflection. These are just elements of the dihedral group corresponding to the symmetries of a regular polygon with vertices corresponding to the nodes of the quivers. Additional symmetries will arise if some of the N_i are equal and we will treat the exchange of groups with identical ranks as giving the same theory. This sort of measure would arise if we imagined our field theory landscape as arising from D-branes on a Calabi-Yau in which all the cycles give rise to gauge theories with the same coupling, which could for example happen if we would resolve an A_k singularity in such a way that all two-cycles would have equal size.

3.2.1 The canonical ensemble breaks down

We will first try to analyze the statistics of cyclic, chiral quivers in a canonical ensemble. All along, as motivated above, we will assume that the gauge groups uniquely fix the matter content. Let r_k be the number of times the group $U(k)$ appears. Then,

the total rank N , and the number gauge factors L , are

$$N = \sum_k k r_k \quad ; \quad L = \sum_k r_k. \quad (20)$$

We want to compute the partition function of this ensemble of *ordered* partitions of N :

$$Z = \sum_{\{r_k\}} \frac{1}{2} \left(\sum_k r_k - 1 \right)! e^{-\beta \sum_k k r_k - \alpha \sum_k r_k} \prod_k \frac{1}{r_k!}. \quad (21)$$

The combinatorial factor that appears here is simply the number of ways we can choose $r_1, r_2, \dots$ gauge group factors out of $\sum_k r_k$, divided by $2(\sum_k r_k)$ to account for the cyclic and reflection symmetry of the quiver³. Rewriting the partition function in terms of the Γ function, we obtain

$$Z = \frac{1}{2} \sum_{\{r_k\}} \Gamma\left(\sum_k r_k\right) \prod_k \frac{e^{-\beta k r_k - \alpha r_k}}{r_k!} \quad (22)$$

Using the integral representation of the Γ function

$$\Gamma(z) = \int_0^\infty dt \, t^{z-1} e^{-t} \quad (23)$$

the partition function can be rewritten as

$$Z = \frac{1}{2} \int_0^\infty dt \, \frac{e^{-t}}{t} \sum_{\{r_k\}} \prod_k \frac{t^{r_k}}{r_k!} e^{-\beta k r_k - \alpha r_k} \quad (24)$$

Exchanging the sum and the product, and after some manipulations, we obtain

$$Z = \frac{1}{2} \int_0^\infty dt \, \frac{e^{-t}}{t} \exp\left(\frac{t e^{-\alpha} e^{-\beta}}{1 - e^{-\beta}}\right). \quad (25)$$

This integral is only convergent if

$$\frac{e^{-\alpha} e^{-\beta}}{1 - e^{-\beta}} < 1 \quad \Rightarrow \quad e^{-\beta} < \frac{1}{1 + e^{-\alpha}} \equiv e^{-\beta_H}. \quad (26)$$

This implies that there is a limiting β above which the partition function is undefined, because the integrand diverges as $t \rightarrow \infty$. There is also always a divergence as $t \rightarrow 0$ which can be regulated by recognizing that the divergence is a constant independent of α and β . To show this, we define $\gamma = \frac{e^{-\alpha} e^{-\beta}}{1 - e^{-\beta}}$, and find that

$$\frac{dZ}{d\gamma} = \int_0^\infty dt \, e^{-(1-\gamma)t} = \frac{1}{1 - \gamma} \quad (27)$$

³This counting actually ignores certain accidental symmetries that can arise in the structure of the quiver. For example, in a cyclic quiver in which the gauge groups $U(N_1)$ and $U(N_2)$ alternate, only one cyclic permutation gives a different configuration for the quiver. The fully correct counting can be derived using Polya theory – we are simply using the leading terms of that analysis, which is sufficient for our purposes.

which implies that, below the limiting temperature,

$$Z = -\log(1 - \gamma) = -\log\left(1 - \frac{e^{-\alpha}e^{-\beta}}{1 - e^{-\beta}}\right) = -\log\left(1 - \frac{uq}{1 - q}\right) \quad (28)$$

where $u = e^{-\alpha}$ and $q = e^{-\beta}$.

In order to achieve a large total rank, β must be tuned to close to its limiting value β_H (26). Then, if, for example, we put $u = 1$, the expectation value of the total rank is

$$\langle N \rangle = q \frac{\partial}{\partial q} \log Z \sim \frac{-1}{2\epsilon \log(4\epsilon)} \quad (29)$$

where we tuned $q = q_H - \epsilon = \frac{1}{2} - \epsilon$ to get a large rank. Similarly, in this approximation we can compute

$$\langle r_k \rangle \sim \left(\frac{1}{2}\right)^{k+1} \frac{-1}{2\epsilon \log(4\epsilon)} \sim \left(\frac{1}{2}\right)^{k+1} \langle N \rangle. \quad (30)$$

This is completely different from the matter-free result for the typical partition: for example, on average one quarter of the nodes will be abelian. However, we also find that

$$\text{Var}(r_k) \sim \left(\frac{1}{2}\right)^{2r+2} \frac{-1}{(2\epsilon)^2 \log(4\epsilon)} \sim -(1 + \log(4\epsilon)) \langle r_k \rangle^2 \quad (31)$$

This is much larger (as $\epsilon \rightarrow 0$) than the expectation value squared. In other words, the number of group factors with a given rank is not typical in the sense of (6).

As in the matter-free case, we might wonder if a more coarse-grained question would have a more statistically predictable answer. For example, we might ask how many gauge factors we expect to see within some range of ranks. The mean and variance of such a coarse grained variable can be extracted by summing over the quantities in (30,31) because the r_k are independent random variables in our ensemble. In the central limit theorem, summing M identically distributed random variables reduces their fluctuations because both the mean and the variance are enhanced by a factor of M ; thus the variance to mean squared ratio is *reduced* by a factor of M . In the matter-free example, something like this happened because, although the r_k were not identically distributed, their dependence on k was sufficiently weak to allow the central limit theorem to work. In the present case, the exponential dependence of (30, 31) on the rank k means that this mechanism fails – the mean and the variance remain dominated by the smallest k in the range of the sum. Thus, it would appear that there is no simple statistically predictable quantity in this landscape.

However, this is in fact happening because the canonical ensemble is breaking down and is not a good approximation of the microcanonical ensemble anymore. The canonical ensemble will reproduce the microcanonical ensemble when the growth of the configuration space with the total rank is slow enough so that, when multiplied by an exponential Boltzmann factor, a nicely localized measure results. Here the Gamma function and the exponential in (22) compete on an equal footing and lead to a widely spread out measure in which the rank of the gauge group fluctuates wildly over the ensemble, leading to a very large variance. Indeed, we should expect this

sort of behavior to occur generally when studying the statistics of quivers since the number of graphs increases rapidly with the number of nodes. Thus we turn to the microcanonical ensemble in order to implement more accurately our constraint on the total rank.

3.2.2 Microcanonical analysis

We consider once more a cyclic quiver and ignore accidental symmetries. The microcanonical partition function for cyclic gauge theories of rank N and L nodes is simply the number of such theories. This is given by the coefficient of q^N in

$$\frac{1}{2L} [q + q^2 + q^3 + \dots]^L. \quad (32)$$

Here the $1/2L$ divides out the cyclic permutations and reflections. We find that

$$Z_L = \frac{1}{2L} \binom{N-1}{N-L}. \quad (33)$$

Summing this over L we can write a partition function which is canonical in the number of nodes and microcanonical in the total rank N :

$$Z(u) = \sum_{L=1}^N u^L Z_L = \frac{(1+u)^N - 1}{2N}. \quad (34)$$

To get the unbiased landscape in which all theories of equal rank have equal weight, we should take $u = 1$, but we will consider other values of u as well. The expectation value of L is

$$\langle L \rangle = u \partial_u \log(Z(u)) = \frac{u(1+u)^{N-1}}{(1+u)^N - 1} N. \quad (35)$$

For the unbiased ensemble with $u = 1$, we get $\langle L \rangle = \frac{N}{2}$ in the large N limit. However, if $u \sim \frac{1}{\sqrt{N}}$, then $\langle L \rangle \sim \sqrt{N}$, and if $u \sim \frac{1}{N}$, then $\langle L \rangle \sim O(1)$. In fact, if $u \sim N^{-a}$, $\langle L \rangle \sim N^{1-a}$. It can be checked that the canonical analysis gives the same expectation values. However, the microcanonical variance in L is

$$\text{Var}(L) = \left(1 - \frac{Nu}{(1+u)^N - 1}\right) \frac{1}{1+u} \langle L \rangle \quad (36)$$

For the three scalings of u , i.e. $u \sim N^{-a}$, the variance in L is some order 1 number times the mean value of L , independent of a . Thus, when $\langle L \rangle$ is large, the variance to mean squared ratio is small, unlike the canonical analysis. This means that in such landscapes the number of gauge factors is typical in the sense of (6) and is therefore highly predictable.

The expectation value for the number of abelian factors is:

$$\langle r_1 \rangle = \frac{1}{Z} \sum_{L=1}^N \frac{u^L}{L} L \binom{N-2L-2}{N-2L-2} = \frac{u^2(1+u)^{N-2}}{(1+u)^N - 1} N \quad (37)$$

When $u = 1$, this becomes $\langle r_1 \rangle = \frac{1}{4}N$ in the large N limit. When $u \sim \frac{1}{\sqrt{N}}$, $\langle r_1 \rangle \sim O(1)$. And when $u \sim \frac{1}{N}$, $\langle r_1 \rangle \rightarrow 0$. In fact, for $u \sim N^{-a}$, $\langle r_1 \rangle \sim N^{1-2a}$. It can be checked that these expectation values match the canonical ensemble. However, the variance in r_1 is much smaller in the microcanonical ensemble. First we compute that

$$\langle r_1^2 \rangle = \frac{1}{Z} \sum_{L=1}^N \frac{u^L}{L} \left\{ L \binom{N-2}{L-2} + L(L-1) \binom{N-3}{L-3} \right\} \quad (38)$$

$$= \frac{u^2(1+u)^{N-4}(u(uN+4)+1)}{(1+u)^N - 1} N \quad (39)$$

Therefore, the ratio of the variance to the mean squared is

$$\frac{\langle r_1^2 \rangle - \langle r_1 \rangle^2}{\langle r_1 \rangle^2} = \frac{1+u \left(4 - \frac{Nu}{(1+u)^N - 1} \right)}{(1+u)^2} \times \frac{1}{\langle r_1 \rangle} \quad (40)$$

The coefficient of $1/\langle r_1 \rangle$ in this expression is of $O(1)$ for $u \sim N^{-a}$, with $0 \leq a \leq 1$.

Pulling everything together, in the unbiased ensemble with $u = 1$, the average number of gauge factors is $N/2$ and the number of abelian factors is $N/4$. Both of these quantities are typical in the sense of (6) and hence highly predictable in this landscape without any coarse-graining. In a biased ensemble with $u \sim 1/\sqrt{N}$, the total number of gauge factors is $O(\sqrt{N})$, and the number of abelian factors is $O(1)$. Since variance is of the same order as the mean for both quantities, the number of gauge factors is typical and thus predictable, but the number of abelian factors is not. In this case, we expect that a coarse-grained statistic, such the fraction of gauge groups in a given range, would be more predictable as in the matter-free case.

Higher Ranks To find the expectation value of the occupation number of rank r , we can insert a “chemical potential” for that rank. So

$$Z(u, \{y_k\}) = \sum_{L=1}^N \frac{u^L}{2L} \left[\sum_{k=1}^N q^k y_k \right]^L \Big|_{q^N}, \quad (41)$$

where the left hand side equals the coefficient of q^N in the right hand side. The expectation value $\langle r_k \rangle$ is given by

$$\langle r_k \rangle = \partial_{y_k} \log \left(Z(u, \{y_k\}) \right) \Big|_{\{y_k\}=1} \quad (42)$$

$$= \frac{1}{Z[u]} \sum_{L=1}^N u^L \binom{N-k-1}{L-2} \quad (43)$$

$$= \frac{u^2(1+u)^{N-k-1}}{(1+u)^N - 1} N \quad (44)$$

In the unbiased ensemble with $u \sim 1$, $\langle r_k \rangle \sim (1/2)^{k+1}N$ as we found in the canonical ensemble. Similarly, the expectation value $\langle N_r^2 \rangle$ is:

$$\langle r_k^2 \rangle = \frac{1}{Z} y_k \partial_{y_k} y_k \partial_{y_k} \left(Z(u, \{y_k\}) \right) |_{\{y_k\}=1} \quad (45)$$

$$= \frac{1}{Z} \sum_{L=1}^N u^L \left\{ \binom{N-r-1}{L-2} + (L-1) \binom{N-2r-1}{L-3} \right\} \quad (46)$$

$$= \frac{u^2(1+u)^{N-2r-2} \left(2u + (N-2r+1)u^2 + (1+u)^{r+1} \right)}{(1+u)^N - 1} N \quad (47)$$

So the ratio of the variance to the mean squared is

$$\frac{\text{Var}(r_k)}{\langle r_k \rangle^2} = \frac{1}{(1+u)^{k+1}} \left\{ (1+u)^{k+1} + u((1-2k)u+2) - \frac{Nu^2}{(1+u)^N - 1} \right\} \times \frac{1}{\langle r_k \rangle} \quad (48)$$

This is always $O(1)$ times $1/\langle r_k \rangle$, and hence the number of gauge groups of a given rank is typical, and hence highly predictable, if the average is large.

Lessons: We are finding that in an anarchic landscape of cyclic quiver gauge theories, the actual number of gauge factors of a given rank is highly predictable. Specifically, the distribution of ranks is exponential and the low rank populations are statistically predictable with high confidence. In a biased landscape in which the measure favors having a number of gauge factors that is sufficiently smaller than the total rank, we found that the number of factors with a fixed rank is not typical in general although the total number of factors can be. In this case, one could test whether an appropriately coarse grained quantity, like the fraction of gauge groups with ranks in some range, is more predictable.

4 Thinking about the general quiver

To extend our analysis to the general quiver gauge theory we could try to compute a partition sum of the form

$$Z = \sum_L \sum_{N_i, A_{ij}} \exp(-\beta \sum_i N_i - \lambda \sum_{ij} A_{ij} N_i N_j) \quad (49)$$

where L is the number of nodes of the quiver, N_i are the ranks of the gauge groups, and A_{ij} are the numbers of bifundamentals between nodes i and j . One difficulty is that this partition sum is canonical and, as we found, it may not implement the constraints on the total rank and the amount of matter very well because of the rapid growth of the space of theories. Secondly the sum should only be over anomaly cancelled theories. Thirdly, there are discrete symmetries which tend to lead to vanishing expectation values. In view of this, below we will develop some approaches to dealing with the two latter issues.

4.1 Implementing anomaly cancellation

A loop basis for anomaly free theories: If all the gauge groups have the same rank, the general anomaly free theory can be constructed by making sure that the bifundamental fields always form closed loops. One can always construct such matter distributions by saying that each of the possible loops in the quiver has n_i fields running around it. Where loops overlap the matter content will either add or subtract depending on the orientation of the loops (again here we are supposing that non-chiral doublets decouple; in addition, we identify negative A_{ij} with a positive A_{ji} and vice versa.). Any loop in the quiver can be constructed by summation of a basis of independent 3-loops and it can be shown that this basis will have

$$N_L = \frac{(L-1)(L-2)}{2} \quad (50)$$

elements. For example, consider the case with $L = 6$ nodes, i.e. there are six gauge groups that we label from 1 to 6. Then, the following three loops form a basis for all loops: (123), (124), (125), (126), (234), (235), (236), (345), (346), (456). The basis has 10 elements which is equal to $N_6 = (6-1)(6-2)/2$. We can check that the N_L loops provide enough free parameters to parameterize the space of anomaly free theories. To see this, note that the solutions to the anomaly cancellation equations form a vector space of dimension

$$\frac{L(L-1)}{2} - (L-1) = \frac{(L-1)(L-2)}{2} = N_L \quad (51)$$

where $L(L-1)/2$ is the number of parameters A_{ij} from which we have subtracted the $(L-1)$ anomaly cancellation conditions on L groups.

Even when the ranks are unequal, anomaly free theories can be constructed from a basis of 3-loops because (50) and (51) are equal. However, the links of any given 3-loop will have to be populated with a different number of fields in a way related to the GCDs of the three groups appearing in it. For example, suppose one has the three gauge groups $SU(r_1 \cdot g) \times SU(r_2 \cdot g) \times SU(r_3 \cdot g)$ where r_i are a triple of positive integers that do not share a common factor and g is any another positive integer. Then if we take number of chiral bifundamentals between gauge group i and j to be $A_{ij} = \epsilon^{ijk} r_k$, we get an anomaly free theory.⁴

This way of thinking suggests that one way to do the statistics of anomaly free theories is to first select a basis of anomaly free 3-loops and then do the statistics of populations of these loops given a bound on the total number of loops.

⁴As a specific example consider a four-node quiver with gauge group $SU(3)_1 \times SU(5)_2 \times SU(7)_3 \times SU(8)_4$. Then, we can get an anomaly free theory by making a loop of four with $A_{12} = 7 \cdot 8$, $A_{23} = 3 \cdot 8$, $A_{34} = 3 \cdot 5$, $A_{41} = 7 \cdot 5$. We can obtain this as a sum of two 3 loops: (124) + (234). The loop (123) corresponds to an $SU(3)_a \times SU(5)_b \times SU(8)_c$ with $A_{ab} = 8$, $A_{bc} = 3$, $A_{ca} = 5$ while the loop (234) corresponds to $SU(5)_i \times SU(7)_j \times SU(8)_k$ with $A_{ij} = 8$, $A_{jk} = 5$, $A_{ki} = 7$. To get the four loop (1234), we need to cancel the (24) link which means that need to add $7 \cdot (124) + 3 \cdot (234)$. This precisely reproduces the four loop numbers. Another anomaly free theory could be generated by adding (124) to (234). In this case, the fields along link (24) will not cancel, but by construction, the number of fields going into and out of each gauge group will cancel.




Figure 1: The five vertices from which all anomaly-free, chiral, asymptotically free, equal rank theories are constructed

Anomaly free, asymptotically free, chiral, equal rank gauge theories:

This set of theories is very easy to analyze, as there can be only five different types of vertices in such quivers (see Fig. 1). Therefore the most general quiver arises by combining these five vertices in various combinations. Superficially, the second vertex with two separate lines coming in and two separate lines going out allows for the largest amount of combinatorial freedom and will quite likely dominate this set of theories. It would be interesting to explore this class further. Possibly it can be mapped to an existing solvable lattice model in statistical mechanics.

Anomaly cancellation for a general quiver by using an extra node:

If we drop the constraint of asymptotic freedom, the set of anomaly free, chiral, and equal rank theories is easy to parametrize. It is not difficult to see that if we take any set of edges $\mathcal{S}$ such that the edges together form a connected tree which contain all vertices of the quiver, then the anomaly equations uniquely determine the A_{ij}, A_{ji} with $(ij) \in \mathcal{S}$ in terms of the A_{ij}, A_{ji} with $(ij) \notin \mathcal{S}$. Thus we can simply take an arbitrary set of chiral matter fields for all edges not in $\mathcal{S}$, after which anomaly cancellation uniquely fixes the remaining links.

A simple example of the set $\mathcal{S}$ is the star-shaped tree consisting of all edges $(1i)$, $i = 2 \dots L$. In other words, if we remove one vertex and all its edges, and arbitrarily specify the chiral matter content in the remaining quiver with $L - 1$ vertices, this uniquely determines an anomaly free, chiral, equal rank quiver gauge theory with L gauge groups. To illustrate this, consider a four-node quiver. Take $A_{12} = a$, $A_{32} = b$ and $A_{13} = c$ ⁵. Then anomaly cancellation uniquely fixes

$$A_{24} = a + b, \quad A_{41} = a + c, \quad A_{43} = b - c. \quad (52)$$

This method can be extended to theories where the gauge groups have unequal ranks. First consider an arbitrary chiral, quiver with $L - 1$ nodes. Let the rank of the group at the i th node be N_i . For anomaly cancellation, the net number of fundamentals minus antifundamentals at each node must be zero. Let K_i be the net excess matter (number of fundamentals minus anti-fundamentals) at each node. Then we can always add an additional $U(1)$ gauge group with $N_i K_i$ bifundamental fields under this $U(1)$ and the $U(N_i)$ of the i th node. This will give an anomaly free theory.

⁵If we say $A_{12} = a$, then we mean that $A_{12} = a$ for $a \geq 0$ and $A_{21} = -a$ for $a \leq 0$. This will guarantee that the theory is chiral

This extra node can be non-abelian, but its rank is restricted to be a divisor of the set $\{N_i K_i\}$. In this way, the statistics of general anomaly free quivers on L nodes can be studied by first constructing arbitrary $L - 1$ node quivers and then adding a extra node according to the above algorithm.

4.2 Dealing with discrete quiver symmetries: an example

From above, the set of anomaly free, chiral and equal rank theories with four nodes is parametrized by the rank N of the gauge groups and three integers a, b, c . The measure (8) becomes

$$\rho = \exp \left(-4\beta N - \lambda N^2 (|a| + |b| + |c| + |a + b| + |a + c| + |b - c|) \right). \quad (53)$$

In the remainder, we will fix the value of N and look only at the distribution of a, b, c .

By symmetry, the expectation values of a, b, c are all zero. This happens because there are a number of discrete symmetries of the quiver due to which averages vanish. For example, for every chiral quiver there is the anti-chiral quiver in which the orientations of all fields are reversed. Averaging these two will formally give $a = b = c = 0$. Similar phenomena will always happen whenever we consider sets of quivers with symmetries. More structure appears once we break the symmetries and look at the average quiver in an ensemble with some symmetry breaking conditions imposed. Suppose for example that we impose $a > 0$. This leaves a $\mathbb{Z}_2$ symmetry that exchanges vertices 3 and 4. Therefore, the expectation value of A_{34} will be zero. Symmetry considerations further show that

$$\langle \frac{1}{2} A_{12} \rangle = \langle A_{23} \rangle = \langle A_{24} \rangle = \langle A_{31} \rangle = \langle A_{41} \rangle. \quad (54)$$

Furthermore, each of these expectation values is proportional to $1/\lambda N^2$.

A boundary condition that completely breaks the symmetry is to impose that $a \geq b \geq 0$. We can always achieve this up to a permutation of the vertices so there is no loss of generality. The analysis of the expectation values of the number of matter fields in this ensemble is more tedious but can still be done explicitly. To leading order in $\epsilon = \lambda N^2$ we obtain⁶

$$\begin{aligned} \langle A_{12} \rangle &= \frac{47}{84\epsilon}, & \langle A_{32} \rangle &= \frac{4}{21\epsilon}, & \langle A_{31} \rangle &= \frac{61}{588\epsilon}, \\ \langle A_{24} \rangle &= \frac{3}{4\epsilon}, & \langle A_{41} \rangle &= \frac{67}{147\epsilon}, & \langle A_{43} \rangle &= \frac{173}{588\epsilon}. \end{aligned} \quad (55)$$

Thus we see that after modding out the $\mathbb{Z}_2$ symmetries of the quiver we are able find an interesting average quiver. Of course, since there are only four nodes here, we do not expect any notion of statistical typicality. To study whether general large quivers have some typical structure, we will have to proceed as above, by parameterizing the space of anomaly cancelled theories and then imposing symmetry breaking conditions.




Figure 2: Examples of flows to the infrared in asymptotically free, four node quiver gauge theories with equal rank gauge groups. The tree level superpotential is assumed to be trivial. (a) This quiver flows to a theory with a Coulomb branch at low energies. (b) This theory is a conformal field theory at low energies. (c) Assuming the two gauge groups with N flavors have a higher scale than the groups with $2N$ flavors, this theory flows to a theory with two confined groups, with the massless mesons of the two groups then participating in an interacting conformal field theory. (d) Assume that the group that confines has a higher dynamical scale than the other groups, and that the confinement is on the Baryonic branch. The massless mesons of this confining factor then interact with the rest of the theory leading to a flow to an interacting conformal field theory.

4.3 Towards dynamics

While we have been focusing on the structure of those field theories in which anomalies cancel, we should also be paying attention to dynamics. Since we are dealing with $\mathcal{N} = 1$ field theories, if $N_f > 3N_c$ for any gauge group then it will be infrared free. If $N_f < 3N_c$ it will be asymptotically free. If $N_f = 3N_c$ the one-loop beta function vanishes. If we distribute fields into a quiver, the bound of the total number of fields will tend to cause the low rank gauge groups to contain more fields. Thus they will tend to be infrared free. What is more, because, as we have seen above, anomaly cancellation including high rank gauge groups tends to require lots of fields, if a high rank group is connected to the rest of the quiver it would tend to push groups in the quiver towards infrared freedom. In general, studying RG flow requires us to know the superpotential or at least to scan statistically over them. Minimally, we should include all cubic and quartic terms in the superpotential with $O(1)$ coefficients multiplied by the appropriate scale. (The cubic terms are classically marginal, and some quartic terms are known to become marginal under RG flow.) Doing such a dynamical analysis of general quiver gauge theories is beyond the scope of this paper, but as an initial step to gain some experience with how this works we will study some examples without a superpotential.

4.3.1 Four node, asymptotically free quivers

First recall that $SU(N)$ gauge theory with N flavors confines at energies below its dynamical scale, while $SU(N)$ theory with $2N$ flavors flows to an interacting conformal fixed point. We will assume that the confining $SU(N)$ theory is on the baryonic branch. We can then naively take a quiver and simply proceed to allow individual gauge factors to confine, Seiberg dualize [21] etc. as their dynamics becomes strong. A cursory analysis of four node, asymptotically free quivers (see some examples with equal ranks N in Fig. 2, constructed from the vertices in Fig. 1) suggests that one will tend to get interacting conformal field theories in which the mesons of the confining factors participate. This suggests that unparticles [22] might be generic in these settings.

4.3.2 General quiver with unequal gauge groups

First consider the simple case of a loop of three gauge groups, $SU(N_1) \times SU(N_2) \times SU(N_3)$ which has to cancel anomalies by itself. For example, this can happen if the 3-loop is isolated within a larger quiver. As we discussed, such primitive 3-loops can be used to generate larger anomaly free quiver gauge theories. To cancel anomalies, the (12), (23), (31) links will generically contain N_3, N_1, N_2 bifundamentals respectively.⁷

⁶Here, by $\langle A_{ij} \rangle$ we really mean $\langle A_{ij} - A_{ji} \rangle$.

⁷The minimal solution to the anomaly cancellation equations will actually be that the number of bifundamentals connecting i and j is $N_k / \text{GCD}(\{N_i, N_j, N_k\})$ as in (19). But generically the GCD will be 1.

Thus for group i to be asymptotically free one will need that

$$3N_i > N_j N_k \quad i \neq j \neq k. \quad (56)$$

Taking all the $N_i > 3$ and $N_1 < N_2 < N_3$, it is clear that $SU(N_3)$ is the only gauge group that has the possibility of being asymptotically free. So for any anomaly-free, chiral connected quiver with three nodes with ranks at least 3 either all three groups are infrared free, or only the largest one is asymptotically free if it has sufficiently large rank.

The same argument no longer works for connected quivers with more than three gauge groups, still it is easy to see that generically high rank gauge groups with links to smaller rank gauge groups have a chance to be asymptotically free, whereas low rank gauge groups connected to higher rank gauge groups tend to be IR free.

Now consider three cases for the dynamics of a quiver with unequal gauge groups.

1. The number of fields K is very large. If so, it seems likely that in a randomly chosen field theory all possible links in the quiver will be populated with some multiplicity, although the links between low rank groups will be enhanced. In this circumstance our arguments suggests that the entire theory will be infrared free.
2. The number of fields K is small. Presumably the lowest rank gauge groups will tend to have matter and the quiver will typically consist of several disconnected smaller clusters that each for a connected quiver gauge theory. The high rank gauge groups with little matter would then confine at the appropriate dynamical scale.
3. For an intermediate number of fields the clusters will percolate and presumably there is an interesting phase structure here.

5 Conclusion

It is somewhat unsettling to attempt to make *statistical* predictions for the structure of the theory describing nature because, ever since Galileo, we have been fortunate that observations and symmetries have constrained the possibilities sufficiently to essentially give a unique theory describing the phenomena under investigation. But string theorists are in the unprecedented situation of hoping to make predictions for the fundamental theory up to the Planck scale given observations below the TeV scale, subject to only very general constraints such as consistency and in particular a consistent coupling to quantum gravity. In such a situation, the best one can do is to predict the likelihood of possible high energy theories, conditioned on the known facts below the TeV scale, the known constraints, and our best guess regarding the measure on the space of theories. This is literally all that we can know. While this sort of Bayesian approach is unfamiliar in particle physics, it is much less unusual in cosmology where one does conceive of ensembles of possible universes or ensembles of

domains with different low-energy physics in a single universe. We would nevertheless like to emphasize that we do not want to exclude the possibility that consistency requirements plus experimental input will eventually yield an (almost) unique fundamental theory, we are merely entertaining the logical possibility that this will turn out to not be the case.

In this paper we have used the uniform measure on specific effective field theory landscapes, but it is not obvious that this is the measure prescribed by string theory. For example, dynamics can play a role in determining the appropriate measure because there can be transitions between vacua with different properties. Also, renormalization group flows can modify the measure in the infrared as theories flow towards their fixed points. Given the correct measure, our analysis could be repeated to find the typical predictions. However, because the uniform measure leads to typicality for some coarse-grained properties, an alternative measure would have to concentrate on an exponentially sparse part of the configuration space in order to change the typical predictions of the uniform measure.

The general approach to model building suggested by these considerations does not involve the usual desert with a high scale GUT. Instead it appears that one would statistically expect a plethora of gauge factors leading to interesting structures at all scales up to the string scale. Amongst these gauge factors there will be some groups with high ranks and others with low ranks. If there is a bound on the total number of matter fields, statistically, the higher rank groups will tend to have fewer fundamentals (since this eats up matter). Thus they will tend towards confinement at a relatively high dynamical scale if all couplings are unified at the string scale. On the other hand if you have too much matter in any group it will tend to infrared triviality. Thus the low rank groups, if they are to have IR dynamics, will tend to be largely decoupled from the high rank groups. Thus if we study the statistics of anarchic landscapes of field theories, conditioned on having interesting low energy dynamics, we will tend towards a structure with dynamical low rank groups largely decoupled from a complex, interacting higher rank sector.

The explicit examples of toy landscapes that we studied in Sec. 3 do not have very interesting dynamics. The matter-free case confines. The ring quivers that we studied in detail are generically infrared free since anomaly cancellation imposes the need for lots of matter unless the individual gauge group ranks conspire to make the GCD in (19) large. Thus we see that conditioning a field theory landscape on having interesting low energy dynamics, along with anomaly cancellation, will be a major constraint, and is likely to significantly modify the measure on the space of theories. It would be amusing if curious number theoretic properties like the appearance of large GCDs will have to be given more weight. It would also be very interesting to explore other measures; for example the results in [14, 15] suggest to weigh rank k gauge group factors with an extra factor of $1/k^2$ compared to the anarchic measures we have been using.

Acknowledgments: We have benefited from conversations with Ofer Aharony, Michael Douglas, Gary Gibbons, Florian Gmeiner, Dieter Lüster, Juan Maldacena, Ya-

sunori Nomura, Carlos Nunez, Al Shapere, Tanmoy Vachaspati, Brian Wecht, and Timo Weigand. We are grateful to the organizers of the Sowers Workshop at Virginia Tech where this work was initiated. V.B. thanks DAMTP, Cambridge, and the Physics Department at Swansea, and A.N. thanks the Physics Departments at Penn, the University of Amsterdam and the IAS for for hospitality during this project. VB was supported by the DOE under grant DE-FG02-95ER40893, by the NSF collaboration grant OISE-0443607, and as the Helen and Martin Chooljian member of the Institute for Advanced Study. AN is supported by a STFC advanced fellowship. JdB is partially supported by the FOM foundation. Finally, we have enjoyed the atriums of the British Library and the British Museum while this paper was being completed.

References

- [1] R. Bousso and J. Polchinski, “Quantization of four-form fluxes and dynamical neutralization of the cosmological constant,” JHEP **0006**, 006 (2000) [arXiv:hep-th/0004134].
- [2] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, “De Sitter vacua in string theory,” Phys. Rev. D **68**, 046005 (2003) [arXiv:hep-th/0301240].
- [3] B. Feldstein, L. J. Hall and T. Watari, ‘Landscape prediction for the Higgs boson and top quark masses,” Phys. Rev. D **74**, 095011 (2006) [arXiv:hep-ph/0608121]; L. J. Hall, M. P. Salem and T. Watari, “Statistical Understanding of Quark and Lepton Masses in Gaussian Landscapes,” Phys. Rev. D **76**, 093001 (2007) [arXiv:0707.3446 [hep-ph]].
- [4] R. Easther and L. McAllister, “Random matrices and the spectrum of N-flation,” JCAP **0605**, 018 (2006) [arXiv:hep-th/0512102].
- [5] N. Arkani-Hamed, S. Dimopoulos and S. Kachru, “Predictive landscapes and new physics at a TeV,” arXiv:hep-th/0501082.
- [6] L. J. Hall, H. Murayama and N. Weiner, “Neutrino mass anarchy,” Phys. Rev. Lett. **84**, 2572 (2000) [arXiv:hep-ph/9911341]; N. Haba and H. Murayama, “Anarchy and hierarchy,” Phys. Rev. D **63**, 053010 (2001) [arXiv:hep-ph/0009174]; A. de Gouvea and H. Murayama, “Statistical test of anarchy,” Phys. Lett. B **573**, 94 (2003) [arXiv:hep-ph/0301050].
- [7] A. E. Nelson and M. J. Strassler, “Suppressing flavor anarchy,” JHEP **0009**, 030 (2000) [arXiv:hep-ph/0006251].
- [8] C. D. Froggatt and H. B. Nielsen, “Hierarchy Of Quark Masses, Cabibbo Angles And CP Violation,” Nucl. Phys. B **147**, 277 (1979);
- [9] G. Gibbons, “Priors”, seminar at conference *The Very Early Universe 25 Years On*, December 2007.

- [10] K. R. Dienes, “Statistics on the heterotic landscape: Gauge groups and cosmological constants of four-dimensional heterotic strings,” *Phys. Rev. D* **73**, 106010 (2006) [arXiv:hep-th/0602286].
- [11] M. R. Douglas, “The statistics of string / M theory vacua,” *JHEP* **0305**, 046 (2003) [arXiv:hep-th/0303194];
- [12] F. Denef and M. R. Douglas, “Distributions of flux vacua,” *JHEP* **0405**, 072 (2004) [arXiv:hep-th/0404116]; J. P. Conlon and F. Quevedo, “On the explicit construction and statistics of Calabi-Yau flux vacua,” *JHEP* **0410**, 039 (2004) [arXiv:hep-th/0409215].
- [13] T. P. T. Dijkstra, L. R. Huiszoon and A. N. Schellekens, “Supersymmetric standard model spectra from RCFT orientifolds,” *Nucl. Phys. B* **710**, 3 (2005) [arXiv:hep-th/0411129].
- [14] R. Blumenhagen, F. Gmeiner, G. Honecker, D. Lust and T. Weigand, “The statistics of supersymmetric D-brane models,” *Nucl. Phys. B* **713**, 83 (2005) [arXiv:hep-th/0411173]; F. Gmeiner, R. Blumenhagen, G. Honecker, D. Lust and T. Weigand, “One in a billion: MSSM-like D-brane statistics,” *JHEP* **0601**, 004 (2006) [arXiv:hep-th/0510170].
- [15] M. R. Douglas and W. Taylor, “The landscape of intersecting brane models,” *JHEP* **0701**, 031 (2007) [arXiv:hep-th/0606109].
- [16] C. Vafa, “The string landscape and the swampland,” arXiv:hep-th/0509212
- [17] H. Ooguri and C. Vafa, “On the geometry of the string landscape and the swampland,” *Nucl. Phys. B* **766**, 21 (2007) [arXiv:hep-th/0605264].
- [18] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, “The string landscape, black holes and gravity as the weakest force,” *JHEP* **0706**, 060 (2007) [arXiv:hep-th/0601001].
- [19] J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev. D* **7**, 2333 (1973); J. D. Bekenstein, “Generalized second law of thermodynamics in black hole physics,” *Phys. Rev. D* **9**, 3292 (1974).
- [20] A.M. Vershik, “Statistical mechanics of combinatorial partitions, and their limit configurations,” *Funkts. Anal. Prilozh.* 30, No.2, 19-30 (1996) (English translation: *Funct. Anal. Appl.* 30, No.2, 90-105 (1996)).
- [21] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl. Phys. B* **435**, 129 (1995) [arXiv:hep-th/9411149].
- [22] H. Georgi, “Unparticle Physics,” *Phys. Rev. Lett.* **98**, 221601 (2007) [arXiv:hep-ph/0703260].